

GREATER BOSTON GASTROENTEROLOGY

- George Dickstein, MD
 - Steven N. Fine, MD
 - Jaime A. Oviedo, MD
 - Mark A. Painter, MD
 - Greta Taitelbaum, MD CM FRCP(C) FACG
- Heidi Peters, RN/C-ANP**
475 Franklin Street, Suite 110
Framingham, MA 01702

Phone: (508) 620-9200

Facsimile: (508) 620-6483

H. PYLORI BREATH TEST

You are scheduled for a H.Pylori Breath Test on:

_____ at: _____

1. **4 weeks prior to test:** you need to be off of all antibiotics and/or Bismuth type medications. Note: Bismuth medications are Pepto-Bismol and all generic forms.

Date to Stop Antibiotics/Bismuth: _____

Date stopped Antibiotics/Bismuth: _____

2. **2 weeks prior to test:** you need to be off of proton pump inhibitors (PPI).

PPI include:	<u>Generic Name</u>	<u>US Brand Name</u>
	Omeprazole	Prilosec
	Lansoprazole	Prevacid
	Rabeprazole	Aciphex
	Sucralfate	Carafate
	Esomeprazole	Nexium
	Pantoprazole	Protonix

Date to Stop PPI: _____

Date stopped PPI: _____

3. **Day of Test:** *Nothing to eat or drink for 6 hours prior to test.*

Note: You can continue to take H2 Blockers, such as Tagamet, Zantac, Axid, and Pepcid and Antacids, such as Maalox, Roloids, Tums, Mylanta, and Gelusic.

It is critical that you follow all of the above instructions for the test to accurately determine if you are infected. ****PLEASE BRING THIS FORM TO YOUR TEST****